

Isolation and characterization of microsatellite markers from the threatened eastern indigo snake (*Drymarchon couperi*)

Brian M. Shamblin · Travis I. Alstad ·
Dirk J. Stevenson · John N. Macey ·
Frankie H. Snow · Campbell J. Nairn

Received: 23 October 2010 / Accepted: 2 November 2010
© Springer Science+Business Media B.V. 2010

Abstract The eastern indigo snake (*Drymarchon couperi*) is the longest snake in North America. Population declines due to extensive loss of habitat led to federal listing as a threatened species in 1978. Knowledge deficiencies regarding the biology of the eastern indigo snake, including population connectivity and gene flow, are impediments to the development of effective conservation and restoration strategies. We describe primers and polymerase chain reaction (PCR) conditions to amplify 22 tetranucleotide and pentanucleotide nuclear microsatellite loci isolated from the eastern indigo snake. We tested primers using 28 shed skins representing 24 individuals from Fort Stewart, Georgia. Primers yielded an average of 4.6 alleles per locus (range 3–7) and an average observed heterozygosity of 0.63 (range 0.46–0.79). These loci should prove useful for individual identification as well as population level analyses of this federally threatened species.

Keywords Eastern indigo snake · Microsatellites · Tetranucleotide · Georgia · *Drymarchon couperi*

The eastern indigo snake (*Drymarchon couperi*) is a federally threatened species native to the Coastal Plain of the southeastern United States. This species has experienced a reduction in its overall range as well as population declines due to habitat loss, diminishing gopher tortoise populations, and gassing of gopher tortoise burrows (US Fish and Wildlife Service 1982). A rangewide phylogeographic survey using mitochondrial DNA markers detected the presence of Gulf and Atlantic clades (Krysko et al. 2010), concordant with a biogeographic pattern described for several taxa of plants and animals (Soltis et al. 2006); however the patterns and scale of population connectivity via natal dispersal remain unresolved and may best be studied using microsatellite markers. Nuclear genetic markers capable of individual resolution may supplement traditional mark-recapture monitoring efforts for this species (Stevenson et al. 2009) as well as prove useful in assessments of population structure. We addressed this need through development of a panel of eastern indigo snake specific microsatellite markers.

Genomic DNA was enriched for microsatellite loci using the methods described by Glenn and Schable (2005). DNA was extracted from two freshly shed skins collected at a site in Coffee County, Georgia using a Qiagen® DNEasy Blood and Tissue Kit. DNA was digested using *RsaI* endonuclease (New England Biolabs) at 37°C for 1.5 h. Double stranded SuperSNX linkers (Glenn and Schable 2005) were ligated to the digested genomic DNA overnight at 16°C. Linker-ligated DNA was recovered using PCR on an Applied Biosystems 9,700 thermal cycler and verified using a 1% agarose gel at 100 V for 1 h.

B. M. Shamblin · T. I. Alstad · C. J. Nairn (✉)
Daniel B. Warnell School of Forestry and Natural Resources,
University of Georgia, Athens, GA 30602, USA
e-mail: jnairn@warnell.uga.edu

D. J. Stevenson
Project Orianne, Indigo Snake Initiative, 414 Club Drive,
Hinesville, GA 31313, USA

J. N. Macey
Directorate of Public Works, Environmental Division,
Wildlife Branch, 1177 Frank Cochran Drive, Building 1145,
Fort Stewart, GA 31314, USA

F. H. Snow
South Georgia College, 100 West College Park Drive,
Douglas, GA 31533, USA

Table 1 Characterization of tetranucleotide and pentanucleotide microsatellite loci from the eastern indigo snake based on 24 shed skins collected at Ft. Stewart, Georgia

| Locus | Primary sequence | Genbank Asc # | Motif | <i>N</i> | <i>A</i> | <i>H_O</i> | <i>H_E</i> | <i>P_{HW}</i> | Range (Bp) |
|--------------|---|---------------|--|----------|----------|----------------------|----------------------|-----------------------|------------|
| <i>Dry05</i> | F: (CAG) ^a CTGCAAGAAAGCCACACATC R: GTTTGAGCGTATGCACCAAGACAA | GU592703 | AATAG ¹³ | 23 | 7 | 0.74 | 0.76 | 0.71 | 232–262 |
| <i>Dry06</i> | F: (CAG)TTCCTACCACCCATCTCCAA R: GTTTGGCAGCAAATGAATTGAACC | GU592704 | AAGG ⁷ | 24 | 6 | 0.75 | 0.75 | 0.41 | 208–232 |
| <i>Dry14</i> | F: (CAG)TGCACCAAAGACAAATTCCTT R: GTTTTCAAGAGATTGCCAAATGGA | GU592705 | CTGTT ⁹ | 24 | 3 | 0.46 | 0.49 | 0.96 | 224–236 |
| <i>Dry24</i> | F: (CAG)GTGAGGAGTGGGTGGAAG R: GTTTACGAACAGGAAGAAGATAGTG | GU592706 | ATCT ¹⁶ | 23 | 5 | 0.78 | 0.79 | 0.02 | 287–303 |
| <i>Dry28</i> | F: GTTTAATGAACATTTGGGCTTTG R: (CAG)ACATCATATTGTCACCGCTAC | GU592707 | AGAT ¹⁰ | 20 | 3 | 0.70 | 0.61 | 0.05 | 425–433 |
| <i>Dry30</i> | F: GTTTGATTAGATTGGCAGGTTGAC R: (CAG)CCATCATCTATCTATCACCATC | GU592709 | AGAT ¹³ | 24 | 4 | 0.62 | 0.64 | 0.67 | 178–190 |
| <i>Dry33</i> | F: (CAG)TATAAGCACTCTATGGCCAA R: GTTTC AATTTCCCTCTTTCTCTT | GU989033 | AGAT ⁹ ... AGAT ⁶ | 24 | 7 | 0.79 | 0.83 | 0.02 | 232–278 |
| <i>Dry35</i> | F: (CAG)AAATAGGCACATAGCAAAAAT R: GTTTAAGCCCAATTCACAGAG | GU592708 | ATCT ¹⁶ | 24 | 6 | 0.71 | 0.73 | 0.51 | 292–312 |
| <i>Dry40</i> | F: (CAG)TATTTCTCCATTATTTACCA R: GTTTTCCCATCTGCTTCTGTTCT | GU592710 | ATCT ¹² | 16 | 4 | 0.50 | 0.67 | 0.18 | 444–456 |
| <i>Dry44</i> | F: (M13) ^b CCCCTGAAAAAAGCTAAAAG R: GTTTGGGAAAAAAGACGATGAGAG | GU592711 | ACTT ¹⁰ | 24 | 3 | 0.50 | 0.50 | 0.08 | 193–201 |
| <i>Dry45</i> | F: (CAG)GAGGGCATGGCAATAAG R: GTTTGGTTTATTATCAGGGGATGT | GU592712 | ATCT ¹³ | 24 | 4 | 0.46 | 0.70 | 0.02 | 191–203 |
| <i>Dry48</i> | F: (CAG)GGTTCTGAGACTATTGGGAG R: GTTTAAATAGCAACCACTTTTCTTC | GU592713 | ACCT ⁷ ...ATCT ¹² | 24 | 4 | 0.67 | 0.64 | 0.15 | 286–314 |
| <i>Dry55</i> | F: GTTTCCCATTTGGATTTCACTTAA R: (M13)CTCCCTTTCCTTTGTCA | GU592714 | ATCT ¹⁶ | 23 | 6 | 0.70 | 0.69 | 0.46 | 343–371 |
| <i>Dry59</i> | F: GTTTGTGCAGATGCTCCAGATG R: (CAG)GGGGTGGTGGACTGAAA | GU592716 | AGAT ⁷ | 24 | 5 | 0.83 | 0.74 | 0.77 | 252–282 |
| <i>Dry58</i> | F: GTTTCCCTGTTGATTCTGGTGAA R: (M13)GGAAAGGATGAGTAAAGTC | GU592715 | AAGG ¹² ...AAAG ¹⁵ | 24 | 5 | 0.79 | 0.71 | 0.91 | 340–360 |
| <i>Dry61</i> | F: (M13)ATAAGCCGCCCTGAGTC R: GTTTGCCTCTGGTTATTCACA | GU592717 | AATC ¹² ...AAAT ⁴ | 24 | 4 | 0.54 | 0.64 | 0.67 | 162–214 |
| <i>Dry63</i> | F: (CAG)GATGAAAATGGGACTCTTAGT R: GTTTAATATTTCCGGCTGTTGTGTA | GU592718 | AGTT ⁹ | 24 | 4 | 0.46 | 0.52 | 0.36 | 196–244 |
| <i>Dry64</i> | F: GTTTCTTTGGATTTGATTTTGTTT R: AATTTCTTTACTCTTCATTGT | GU592719 | ATCT ¹³ | 24 | 4 | 0.46 | 0.65 | 0.03 | 295–307 |
| <i>Dry65</i> | F: (CAG)AGGGGGTAGTCCTCATCTC R: GTTTCAGGGCTTTGGACAATATG | GU592720 | ATCT ¹⁴ | 23 | 7 | 0.61 | 0.72 | 0.62 | 164–202 |
| <i>Dry68</i> | F: (CAG)TTTGAGTCTTATTGGATTGAGT R: GTTTGAGATGGTGCTGATGAAAT | GU592721 | ACAT ¹⁰ | 24 | 4 | 0.58 | 0.57 | 0.87 | 205–233 |
| <i>Dry69</i> | F: (CAG)ATGTGTGAACCAAATCCTAAT R: GTTTCACGTATTTGTTTTCTCCT | GU592722 | ATGT ⁹ | 24 | 4 | 0.54 | 0.60 | 0.08 | 428–452 |
| <i>Dry70</i> | F: (CAG)GAGGATTAGGGGATTTTTACA R: GTTTAAAGCAAGGGCCAGTAAG | GU989034 | GGAT ⁴ ... AGAT ⁹ | 24 | 3 | 0.58 | 0.51 | 0.76 | 258–266 |

F Forward primer sequence, *R* reverse primer sequence, *N* number of individuals successfully genotyped; *A*, number of alleles, *H_O* observed heterozygosity, *H_E* expected heterozygosity, *P_{HW}* probability of Hardy–Weinberg disequilibrium, *Range* amplicon size in base pairs

^a CAG tag sequence is CAGTCGGGCGTCATCA

^b M13 tag sequence is GGAAACAGCTATGACCAT

Recovered DNA was hybridized to two different mixtures of biotinylated oligonucleotide probes (Integrated DNA Technologies, Mix 1: (AAAC)₆, (AAAG)₆, (AATC)₆, (AATG)₆, (ACAG)₆, (ACCT)₆, (ACTG)₆; Mix 2: (AAAT)₈, (AACT)₈, (AAGT)₈, (ACAT)₈, (AGAT)₈). Probes and hybridized DNA were captured using streptavidin coated magnetic beads (DynaBeads[®], Invitrogen[®]). Enriched DNA was washed and eluted at 95°C, and the enrichment procedure was repeated to increase the efficiency of microsatellite array capture. Microsatellite enriched DNA was ligated into a pCR[®]2.1-TOPO[®] vector and transformed into TOP 10[®] competent cells (Invitrogen).

A total of 384 colonies were screened, treated with ExoSAP-IT[®] (New England Biolabs) and sequenced using BigDye v3.1 (Applied Biosystems). Sequencing reactions were purified using Sephadex G-50 Fine (Sigma–Aldrich) and analyzed on an Applied Biosystems 3730xl DNA Analyzer. Sequences were edited using SeqMan[®] (Lasergene). Contigs were searched for microsatellite loci using MSATCOMMANDER (Faircloth 2008). We designed primers for 72 loci using Primer3.0 (Rozen and Skaletsky 2000). Amplification was performed in 10 µl reactions consisting of 10 mM Tris pH 8.4, 50 mM KCl, 0.5 µM GTTT “pig-tailed” locus specific primer, 0.05 µM CAG (CAGTCGGGCGTCATCA) or M13 (GGAAACAGCTA TGACCAT) tagged locus specific primer, 0.45 µM fluorescently labeled CAG or M13 tag (Boutin-Ganache et al. 2001), 1.5 mM MgCl₂, 0.125 mM dNTPs, 0.5 U Taq polymerase, and approximately 10 ng of template DNA. Fluorophores used were VIC, PET, NED (Applied Biosystems) and FAM (Integrated DNA Technologies). Touchdown cycling parameters (Don et al. 1991) were 95°C for 5 min, 20 cycles of 95°C for 30 s, 60°C minus 0.5°C per cycle for 30 s and 72°C for 1 min, followed by 24 cycles of 95 °C for 30 s, 50°C for 30 s, 72°C for 1 min and a final extension of 72°C for 10 min. Amplicon sizes were analyzed on an Applied Biosystems 3730xl DNA Analyzer using LIZ500 size standard (Applied Biosystems). Allele sizes were scored in GENEMAPPER (Applied Biosystems).

Seventy-two primer pairs were screened for amplification consistency and polymorphism on 28 shed skins from Fort Stewart, Georgia, USA. Twenty-two loci amplified consistently and were polymorphic (Table 1). Data were exported from GENEMAPPER using GMCONVERT (Faircloth 2006), and CERVUS 3.0 (Kalinowski et al. 2007) was used to calculate allelic richness (*A*), observed heterozygosity (*H_O*), and expected heterozygosity (*H_E*). We calculated linkage disequilibrium (LD) and deviations from Hardy–Weinberg equilibrium (HWE) using GENEPOP (Raymond and Rousset 1995). No loci deviated significantly from HWE following sequential Bonferroni correction (table-wide alpha = 0.05; Rice 1989). No pairwise loci comparisons

exhibited linkage disequilibrium following sequential Bonferroni correction. The number of alleles per locus ranged from 3 to 7, with an average of 4.6 alleles per locus.

Three multi-locus genotypes were recovered more than once in the data set. Given the non-exclusion probability of identity of 2.74×10^{-17} with the combined panel, there is sufficient power to infer that these shed skins represent multiple sampling of the same individual through time. These data clearly demonstrate the utility of this panel in discerning individuals for genetic mark-recapture studies. These loci should also prove useful in addressing population level questions in this threatened species.

Acknowledgments For help in collecting specimens and other assistance we thank L. Carlile, M. Hodges, R. Moulis, R. Redmond, A. Safer, R. Tate, M. Wallace, N. Hyslop, J. Waters, and C. Jenkins. We thank D. Alessandrini and the Greater Cincinnati Herpetological Society for a grant to DJS. Tissue samples were handled under the University of Georgia Animal Use Permit #A2010 4-063. Specimens were collected under Georgia Department of Natural Resources Scientific Collection Permit 29-WCH-07-89. We gratefully acknowledge funding provided by the University of Georgia Warnell School of Forestry and Natural Resources.

References

- Boutin-Ganache I, Raposo M, Raymond M, Deschepper CF (2001) M13-tailed primers improve the readability and usability of microsatellite analyses performed with two different allele sizing methods. *Biotechniques* 31:24–28
- Don RH, Cox PT, Wainwright BJ (1991) Touchdown PCR to prevent spurious priming during gene amplification. *Nucleic Acids Res* 19:4008
- Faircloth BC (2006) GMCONVERT: file conversion for Genemapper output files. *Mol Ecol Notes* 6:968–970
- Faircloth BC (2008) MSATCOMMANDER: detection of microsatellite repeat arrays and automated, locus-specific primer design. *Mol Ecol Resour* 8:92–94
- Glenn TC, Schable NA (2005) Isolating microsatellite DNA loci. In: Zimmer EA, Roalson EH (eds) *Methods in enzymology* 395, molecular evolution: producing the biochemical data, part B. Academic Press, San Diego, CA
- Kalinowski ST, Taper ML, Marshall TC (2007) Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Mol Ecol* 16: 1099–10060
- Krysko KL, Smith DJ, Smith CA (2010) Historic and current distribution and preliminary evidence of population genetic structure in the eastern indigo snake (*Drymarchon couperi*) in the southeastern United States. Unpublished report to Florida Fish and Wildlife Conservation Commission
- Raymond M, Rousset F (1995) GENEPOP (version 1.2): population genetics software for exact tests and ecumenicism. *J Heredity* 86:248–249
- Rice WR (1989) Analyzing tables of statistical tests. *Evolution* 43: 223–225
- Rozen S, Skaletsky HJ (2000) PRIMER3 on the WWW for general users and for biologist programmers. In: Krawetz S, Misener S (eds) *Bioinformatics methods and protocols: methods in molecular biology*. Humana Press, Totowa, NJ, pp 365–386

-
- Soltis DE, Morris AB, Machlan JS, Manos PS, Soltis PS (2006) Comparative phylogeography of unglaciated eastern North America. *Mol Ecol* 15:4261–4293
- Stevenson DJ, Enge KM, Carlile LD, Dyer KJ, Norton TM, Hyslop NL, Kiltie RA (2009) An eastern indigo snake (*Drymarchon couperi*) mark-recapture study in southeastern Georgia. *Herpetol Conserv Biol* 4:30–42
- US Fish and Wildlife Service (1982) Eastern Indigo snake recovery plan. U.S. Fish and Wildlife Service, Atlanta, Georgia, p 23